

AASHA PEDIATRICS, INC.

Meena Sathappan, MD
1569 Lexann Avenue #230
San Jose, CA 95121
Ph (408) 274-9099
Fx (408) 274-9009

AUTHORIZATION FOR RELEASE OF MEDICAL RECORDS

TO: _____

Doctor/Office Name

Address

(____) _____

Phone Number

(____) _____

Fax Number

Please **FAX/MAIL** copies of the following information to Aasha Pediatrics, Inc:

Hospital Chart

X-ray Reports

Birth Records

Laboratory Reports

Office Chart

Vaccine Records

Records in the chart from previous physician / facilities

Other: _____

Regarding Patient Name

Date of Birth

Signed: _____

Date: _____

Printed Name: _____

Phone# _____

If not signed by patient, please indicate relationship: Parent/guardian of minor patient
 Personal representative of minor patient

Authorization valid for 90 days only, and may be revoked in writing at any time prior to 90 days by notifying the office.

Meena Sathappan, MD
1569 Lexann Ave, # 230
San Jose, CA 95121

Tel: (408) 274-9099
Fax: (408) 274-9009
www.aashapediatrics.com

PERMIT FOR EAR PIERCING (with Informed Consent)

Informed Consent

Child's Name _____

Address _____

City _____ State _____ Zip _____

Phone (____) _____ - _____

D.O.B of customer ____ / ____ / ____

If your child is under 3 months of age, your infant must have their first DPT inoculation.

I understand that my child's ears will be pierced with pre-sterilized, single-use ear piercing studs that are packaged in a sealed container.

I, the undersigned, acknowledge that I am aware that the ear piercing may carry some risks. These risk include, but are not limited to infection, metal sensitivity, allergic reaction, inflammation, embedding, scarring, fainting and other complications.

I have read, and understand the AFTER CARE PROCEDURES and have received a copy for my reference.

I have agreed to this ear piercing procedure for my child, and I am fully aware of the potential risks and complications.

Parent/ Legal Guardian Signature

____ / ____ / ____

Print Name

Aasha Pediatrics

Meena Sathappan, MD

Lexann Professional Center

1569 Lexann Ave. #230

San Jose, CA 95121

Phone: (408) 274-9099

Fax: (408) 274-9009

HOME CARE INSTRUCTIONS AFTER EAR PIERCING

- Thoroughly cleanse hands with an antibacterial soap prior to any contact with your child's newly pierced ears. DO NOT remove the studs for six weeks.
- Apply antibiotic ointment (Neosporin or Bacitracin) to the front and back of the ear lobe 3 times daily. Gently slide the studs back and forth in your child's ear so that the antibiotic will go inside the piercing. Gently rotate the studs (like winding a watch) 2 to 3 times a day.
- After 6 weeks, the studs can be removed and other post type earrings can be inserted in your child's ears. Post earrings must be worn at all times for the entire first 6 months, to ensure that your child's ear piercing will not close.

If you have any questions, contact us at the phone number above.

Aasha Pediatrics

Meena Sathappan, MD

Lexann Professional Center

1569 Lexann Ave. #230

San Jose, CA 9512

Phone: (408) 274-9099

Fax: (408) 274-9009

HOME CARE AFTER CIRCUMCISION

1. Gently clean the diaper area with only warm water for 24 hours. After that you may return to using baby wipes or your usual method of cleaning.
2. Use an antibacterial ointment, e.g. Bacitracin or Neosporin on a gauze with every diaper change. Apply to the cut edge of the circumcision for the next 7 days to help prevent infections and to keep the edge from sticking to the diaper.
3. There is gauze wrapped around the area of circumcision. The gauze needs to be removed the next day. Soaking it with warm water and waiting 10 minutes before removing the gauze will make the gauze easier to remove.
4. Some oozing of blood from the area of the circumcision is common for the first 24 hours. This requires no special care. However, if the area is bleeding freely, apply pressure to the bleeding area with a clean gauze or cloth and call the office immediately.
5. Your baby can eat as usual after the circumcision.
6. The area of circumcision will take about two weeks to completely heal. During this time you may see whitish or yellowish tissue on the head of the penis. This is not pus and is a part of the normal healing process. However, if you notice a pus-like discharge (usually wipes off easily), please call the office.
7. If the area of the circumcision becomes very swollen, or if your child has a fever (above 100.4) call the office immediately.
8. After 2 weeks, you should begin pulling back the remainder of the foreskin gently for cleaning once a day. This will help ensure that no adhesions develop.
9. Infant Acetaminophen (Tylenol) can be given orally every 4 to 6 hours, for the first 24 hours if needed. (no more than 5x in 24 hours)
10. Follow up in 1 week for circumcision re-check.

I have received and understood these instructions.

Signature

Relationship

Date